

From Local Regge Calculus towards Spin Foam Formalism?

Gabriele Gionti S.J.

Vatican Observatory Research Group,

Steward Observatory, 933 North Cherry Avenue,

The University of Arizona, Tucson AZ 85721 U.S.A. and

*Specola Vaticana, V-00120 Città Del Vaticano, Vatican City State**

Abstract

We introduce the basic elements of $SO(n)$ -local theory of Regge Calculus. A first order formalism, in the sense of Palatini, is defined on the metric-dual Voronoi complex of a simplicial complex. The Quantum Measure exhibits an expansion, in four dimensions, in characters of irreducible representation of $SO(4)$ which has close resemblance and differences as well with the Spin Foam Formalism. The coupling with fermionic matter is easily introduced which could have consequences for the Spin Foam Formalism and Loop Quantum Gravity.

PACS numbers: 04.60.Nc, 04.60.Pp

Keywords: Local Regge Calculus, Loop Quantum Gravity, Spin Foam

*Electronic address: ggionti@as.arizona.edu

I. INTRODUCTION

Regge Calculus [1] (see also [2] for a recent and updated summary of Regge Calculus and its alternative approaches) provides a discretized version of General Relativity. It is, mainly, based on the idea of substituting a Piecewise-Linear (PL)-manifold for differential manifold. In particular, these are simplicial manifolds built by gluing together two distinct n -dimensional flat simplexes by one and only one $(n - 1)$ -dimensional face. The final product of this construction is called a simplicial complex, which owns the manifold structure if it has been made in such a way that each point of the simplicial complex has a neighbourhood homeomorphic to $\mathbb{R}^n$ [3]. The simplicial manifolds, we consider, are orientable. On a simplicial manifold we can define a metric structure. The metric is Euclidean almost everywhere except near the $(n - 2)$ -simplexes called hinges h . Near the hinges the metric is equivalent to a *cone*-metric, that is the metric $\mathbf{g}$ of $\mathbb{R}^{n-2} \times$ the two dimensional cone [4] [1]. A two-dimensional cone is formed by cutting away a circular sector of angle θ and gluing together the two free edges. Its apex is called cone point of angle θ and of curvature $2\pi - \theta$. On each $(n - 2)$ -dimensional simplex h , called an *hinge*, a deficit angle $K(h)$, which is the cone curvature, is defined

$$K(h) = 2\pi - \sum_{\sigma_h^i \supset h} \theta(\sigma_h^i, h) \quad (1)$$

where σ_h^i , $i = 1, \dots, p$ is one of the p n -dimensional simplices incident on h , and $\theta(\sigma_h^i, h)$ its dihedral angles on h . The dihedral angle of a n -simplex on the hinge is the angle between the $(n - 1)$ -dimensional faces that have the hinge in common. $K(h)$ and the volume of the hinge $V(h)$ can be expressed [5] as functions of the squared length of the one-dimensional simplices (edges) of the complex. The Einstein-Regge action is, in analogy to the continuum case, a functional over the simplicial manifolds and depends on the incidence matrix of the simplicial complex [10] and on the (squared) lengths of the edges. Usually, in Regge calculus, the incidence matrix is fixed, so that the action can be written as

$$S_R = \sum_h K(h)V(h) \quad . \quad (2)$$

The corresponding Einstein equations [1] are derived by requiring that the action is stationary under the variation of the length of the edges.




FIG. 1: Dual Voronoi Plaquette

The original aim of this theory was to give approximate solutions of the Einstein equations in the case in which the topology is known. The theory is, as stressed by Regge himself, completely coordinate independent.

Now we summarize and update some definitions and results of [6] that will be useful for our future discussions. Consider an hinge h and suppose that it is shared by $k + 2$ n -simplices (see FIG.1) $\{\alpha, \beta, \delta_1^h, \dots, \delta_k^h\}$.

Given a n -dimensional simplicial complex, there exists a general procedure for defining the dual metric complex, called the Voronoi dual [6] [7]. The n -dimensional Voronoi polyhedron dual to a vertex P of the simplicial complex is the set of the points of the simplicial complex closer to P than to any other vertex, using the standard flat metric in the simplex. It turns out that the k -simplex of the simplicial complex is dual to a $(n - k)$ -polyhedron in the dual Voronoi complex, and the k -dimensional linear space identified by the k simplex is orthogonal to the $(n - k)$ -dimensional space spanned by the corresponding polyhedron. In particular the point dual to the n -simplex S is the centre of a $(d - 1)$ -dimensional sphere passing through all vertices of S (look at FIG.1).

It was pointed out in reference [6] that a theory of Regge Calculus locally invariant under the Poincaré group can be formulated by choosing in every simplex an orthonormal reference frame. In this way every n -dimensional simplex, considered as a piece of $\mathfrak{R}^n$, can be seen

as local inertial reference frame, being flat, and an n -bein base can be chosen in it. As in the continuum theory, we need a connection matrix between the reference frames of two simplexes that share a common $(n - 1)$ -dimensional simplex. Fixing [6] [8] [9] the origin of the reference frames on the dual Voronoi vertices, the orthogonal matrix $\Lambda_b^a(\alpha, \beta)$, which allows passing from the reference frame of α to the β 's, happens to be the Levi-Civita [10], Torsion-free, connection matrix which is a function of the vertex coordinates of the common face $\alpha\beta$ both in the reference frame of α and β .

The hinge h is in one-to-one correspondence with its dual two-dimensional Voronoi plaquette that we still label by h . Now consider the following holonomy matrix and plaquette variable

$$W_\alpha(h) = \Lambda(\alpha, \beta)\Lambda(\beta, \delta_1^h)\dots\Lambda(\delta_k^h, \alpha) \quad . \quad (3)$$

As it has been shown [5] [8], (3) is a rotation in the two-dimensional plane orthogonal to the $(n - 2)$ -dimensional hyperplane spanned by the hinge h . The rotation angle is the deficit angle (1).

We consider $n - 2$ linear independent edge-vectors of the hinge h defined in the following way

$$E_1^a(\alpha) \equiv y_1^a(\alpha) - y_{n-1}^a(\alpha), \dots, E_{n-2}^a(\alpha) \equiv y_{n-2}^a(\alpha) - y_{n-1}^a(\alpha), \quad (4)$$

and the antisymmetric tensor related to the oriented volume of h

$$\mathcal{V}^{(h) \ ab}(\alpha) \equiv \frac{1}{(n - 2)!} \epsilon_{c_1 \dots c_{n-2}}^{ab} E_1^{c_1}(\alpha) \dots E_{n-2}^{c_{n-2}}(\alpha) \quad . \quad (5)$$

At this point it seems natural to propose the following gravitational action [6]

$$I = -\frac{1}{2} \sum_h \left(W_\alpha^{a_1 a_2 \ (h)} \mathcal{V}^{(h)}_{a_1 a_2}(\alpha) \right) \quad . \quad (6)$$

Let $V(h)$ be the oriented volume of the hinge h . The action (6) is equal [6] to

$$I = \sum_h \sin K(h) V(h) \quad , \quad (7)$$

which for small deficit angles $K(h)$ reduces to the Regge action (2).

The presence in the action of $\sin K(h)$ instead of $K(h)$ is only a lattice artifact [6]. The useful regime of this theory is $\sin K(h) \approx K(h)$. In fact Fröhlich [10] has conjectured, as

claimed also in [6], that this action converges to the Einstein-Hilbert action, and J.Cheeger, W. Muller, R.Schrader in [11] have proved that the Einstein-Regge action (2) converges to the Einstein-Hilbert action. They proved that the convergence of the action is in *sense of measure*. This applies when, roughly speaking, the number of the hinges of the simplicial manifold increases along with the incident number of the simplices at each hinge. Then the triangulations will be finer and finer and the difference, in modulus, between the Einstein-Hilbert action on a manifold and the Regge-Einstein one, on the triangulations of the same manifold, become smaller and smaller.

II. FIRST-ORDER FORMALISM

Let's consider the $(n - 1)$ -dimensional face $f_{\alpha\beta} \equiv \{P_1, \dots, Q_{\alpha\beta}\}$ between the simplices α and β . The vertex coordinates of $f_{\alpha\beta}$ are $x_i^a(\alpha)$, $a = 1, \dots, n$. The following vector [6]

$$\begin{aligned} b_{\alpha\beta}^a(\alpha) &= \epsilon_{ab_1 \dots b_{n-1}} (x_1(\alpha) - x_n(\alpha))^{b_1} \dots (x_{n-1}(\alpha) - x_n(\alpha))^{b_{n-1}} \\ &= \epsilon_{ab_1 \dots b_{n-1}} E_1^{b_1} \dots E_{n-1}^{b_{n-1}} \end{aligned} \quad (8)$$

is normal to the face $f_{\alpha\beta}$, and it is assumed to point outward from the interior of the simplex α . The analogous vector $b_{\beta\alpha}^a(\beta)$ in the reference frame of β is related to the previous one by

$$b_{\alpha\beta}^a(\alpha) = \Lambda_b^a(\alpha, \beta) b_{\beta\alpha}^b(\beta) \quad . \quad (9)$$

It can be easily proved that (5) can be written as a bivector

$$\mathcal{V}^{(h) \ c_1 c_2}(\alpha) \equiv \frac{1}{n!(n-2)!V(\alpha)} \left(b_{\alpha\beta}^{c_1}(\alpha) b_{\alpha\delta_k^h}^{c_2}(\alpha) - b_{\alpha\beta}^{c_2}(\alpha) b_{\alpha,\delta_k^h}^{c_1}(\alpha) \right), \quad (10)$$

where $V(\alpha)$ is the oriented volume of the simplex α .

We can consider the action as written on the dual Voronoi-complex of the original simplicial complex. The dual of the hinge h is a two dimensional plaquette whose vertices will be $\{\alpha, \beta, \delta_1^h, \dots, \delta_k^h\}$. These vertices are the dual of the n -simplices incident on the hinge h . If, briefly, we indicate $\Lambda(\alpha, \beta)$ as $\Lambda_{\alpha\beta}$, and so on, the holonomy matrix (3) around the

plaquette is

$$U_{\alpha\alpha}^h \equiv \Lambda_{\alpha\beta}\Lambda_{\beta\delta_1^h}\dots\Lambda_{\delta_k^h\alpha} \quad , \quad (11)$$

However this is still a second order formalism of discrete General Relativity . The connection matrices $\Lambda_{\alpha\beta}$ and the $b_{\alpha\beta}$ are both functions of the coordinates of the edges of the simplicial complex. Now we introduce a first order formalism in which $\Lambda_{\alpha\beta}$ and $b_{\alpha\beta}(\alpha)$ are independent variables. Then equation (9) becomes a constraint

On the $b_{\alpha\beta}(\alpha)$ there is a further constraint since the $n+1$ normals to the $(n-1)$ -dimensional faces of a n -simplex α are linearly dependent,

$$\sum_{\beta=1}^{n+1} b_{\alpha\beta}(\alpha) = 0 \quad . \quad (12)$$

There is a technical problem we would like pointing out. The gravitational action in the form (6) could be dependent on the starting simplex . So we need to define the following antisymmetric tensor on the plaquette h

$$\begin{aligned} W_{c_1 c_2}^{(h)}(\alpha) \equiv & \frac{1}{k_h + 2} \left(\mathcal{V}^{(h)}(\alpha) + \Lambda_{\alpha\beta} \mathcal{V}^{(h)}(\beta) \Lambda_{\beta\alpha} + \dots \right. \\ & \left. + \Lambda_{\alpha\beta} \dots \Lambda_{\delta_{k-1}^h \delta_k^h} \mathcal{V}^{(h)}(\delta_k^h) \Lambda_{\delta_k^h \delta_{k-1}^h} \dots \Lambda_{\beta\alpha} \right)_{c_1 c_2} . \end{aligned} \quad (13)$$

The action then is

$$S \equiv -\frac{1}{2} \sum_h \text{Tr} (U_{\alpha\alpha}^h W^h(\alpha)) \quad (14)$$

The action (14) coincides with the action (6) in the second-order formalism, and, since (in matrix notation)

$$W^{(h)}(\alpha) = \Lambda_{\alpha\beta} \dots \Lambda_{\delta_{i-1} \delta_i} W^{(h)}(\delta_i) \Lambda_{\delta_i \delta_{i-1}} \dots \Lambda_{\beta\alpha} \quad , \quad (15)$$

equation (14) is independent of the starting simplex.

Moreover the action (14) is invariant under the following set of transformations

$$\begin{aligned} \Lambda_{\alpha\beta} &\mapsto \Lambda'_{\alpha\beta} = O(\alpha) \Lambda_{\alpha\beta} O^{-1}(\beta) \\ b_{\alpha\beta}(\alpha) &\mapsto b'_{\alpha\beta}(\alpha) = O(\alpha) b_{\alpha\beta}(\alpha) \end{aligned} \quad (16)$$

where $O(\alpha)$ and $O(\beta)$ are two elements of $SO(n)$. This can be interpreted as the diffeomorphism invariance of this discrete theory of Gravity.

III. FROM QUANTUM MEASURE TO SPIN FOAM

In this section we shall mainly focus on the four dimensional case. The action (14) is invariant under the group $SO(4)$ which is the gauge group. It has been proved [8] [9] that the discrete equations of motion for this theory can be derived and appear very similar to General Relativity equations in the covariant tetrad formalism [12], and in the limit for *small deficit angles* the *Levi-Civita-Regge* connection (the second order formalism which, in this case, is the Regge Calculus itself) is, locally, the unique solution. This is, clearly, an analogy with the Palatini continuum theory where, in the Torsion-free case, the Levi-Civita connection is solution of one part of the first order field equations.

We will use the following notation:

$$\mu(b_{\alpha\beta}(\alpha)) \equiv db_{\alpha\beta}^1(\alpha) \dots db_{\alpha\beta}^n(\alpha) \quad , \quad (17)$$

and let

$$\mu(\Lambda_{\alpha\beta}) \quad (18)$$

be the Haar measure on $SO(4)$. The partition function for this theory is:

$$Z = \int e^{\frac{1}{2} \sum_h \text{Tr}(U_{\alpha\alpha}^h W^h(\alpha))} \prod_{\alpha} \delta\left(\sum_{\beta=1}^{n+1} b_{\alpha\beta}(\alpha)\right) \prod_{\alpha\beta} \delta(b_{\alpha\beta}(\alpha) - \Lambda_{\alpha\beta} b_{\beta\alpha}(\beta)) \mu(\Lambda_{\alpha\beta}) \mu(b_{\alpha\beta}) \quad . \quad (19)$$

And this path-integral is invariant [9] under local $SO(4)$ transformations (16).

From Group Theory [13] it is well known that a function f on a compact group G is called central if $\forall g, h \in G, f(h^{-1}gh) = f(g)$. Thanks to the gauge invariance (16) previously mentioned $e^{\frac{1}{2} \text{Tr}(U_{\alpha\alpha}^h W^h(\alpha))}$ is a central function on $SO(4)$. It follows [13] we can expand it in terms of the characters $\chi_J(U^h)$ of the irreducible representations J of the group $SO(4)$. We can omit the index $\alpha\alpha$ in the holonomy matrix U^h since it does not depend from the

starting simplex α .

$$e^{\frac{1}{2}Tr(U_{\alpha\alpha}^h W^h(\alpha))} = \sum_J c_J^h(b_{\alpha\beta}, b_{\beta\alpha}, \dots, b_{\delta_k^h \alpha}, b_{\alpha \delta_k^h}) \chi_J(U^h) \quad , \quad (20)$$

where the coefficients $c_J^h(b_{\alpha\beta}, b_{\beta\alpha}, \dots, b_{\delta_k^h \alpha}, b_{\alpha \delta_k^h})$ are defined by [13]

$$c_J^h(b_{\alpha\beta}, b_{\beta\alpha}, \dots, b_{\delta_k^h \alpha}, b_{\alpha \delta_k^h}) \equiv \int_{SO(4)} \mu(U^h) e^{\frac{1}{2}Tr(U_{\alpha\alpha}^h W^h(\alpha))} \overline{\chi_J(U^h)} \quad , \quad (21)$$

$\mu(U^h)$ being the $SO(4)$ Haar measure on the holonomy matrices (U^h) and $\overline{\chi_J(U^h)}$ the complex conjugate of $\chi_J(U^h)$.

These new considerations imply we can write the previous Quantum Measure as

$$Z = \int \prod_h \sum_J c_J^h(b_{\alpha\beta}, b_{\beta\alpha}, \dots, b_{\delta_k^h \alpha}, b_{\alpha \delta_k^h}) \chi_J(U^h) \prod_{\alpha} \delta\left(\sum_{\beta=1}^{n+1} b_{\alpha\beta}(\alpha)\right) \prod_{\alpha\beta} \delta(b_{\alpha\beta}(\alpha) - \Lambda_{\alpha\beta} b_{\beta\alpha}(\beta)) \mu(\Lambda_{\alpha\beta}) \mu(b_{\alpha\beta}) \quad , \quad (22)$$

and this last equation seems suggesting that we can label the interior of each plaquette by an irreducible representation J of $SO(4)$. This is very similar to the Spin Foam Formalism where the interior of each oriented 2-polyhedron is labeled, or colored, by an irreducible representation of $SO(4)$ [14][15]. Although this is an early first stage result some sensible differences appear respect to Spin Foam Formalism. The main one, we would like to highlight, is that Barret and Crane [16] proved that the geometry of each four-simplex can be completely characterized by ten bivector two-forms

$$B_{IJ} \equiv E_I \wedge E_J \quad (23)$$

where E_I and E_J are two independent edge vectors which belong to one of ten triangles of a four-dimensional simplex. In order to label each oriented 2-polyhedron by an irreducible representation of $SO(4)$ they quantize the bivector two-form. Then they associate, in four dimensions, an irreducible representation J to the interior of each triangle. Some technical problems arise in implementing this quantization[17]. Mainly in the simplicial BF theory for Spin Foam Formalism, which is the proposal of Barret-Crane model[16], it is needed to consider a generic two form B_{IJ} and impose, at quantum level, that equation (23) be a quantum constraint.

Although we have also a bivector two-form W^h (13) in the action (14), which has a close resemblance with BF and Plebanski's action for General Relativity, it is very clearly defined on the 2- Voronoi-plaquette instead of the triangle dual to the plaquette. Moreover we do not need to quantize this bivector two-form because, as we have seen, the expansion (20) in the path-integral tells us how to label the plaquette by an irreducible representation of $SO(4)$.

Anyway, till now, we have not found that the partition function (22) can be written as claimed in Spin Foam Formalism [14]:

$$Z = \sum_{\Gamma} w(\Gamma) \sum_{J_f, i_e} \prod_f A_f(j_f) \prod_e A_e(j_f, i_e) \prod_v A_v(j_f, i_e) \quad . \quad (24)$$

where Γ is an oriented 2-complex, $w(\Gamma)$ is a weight associated to each 2-complex Γ , J_f is an $SO(4)$ -irreducible representation associated to each 2-face f , i_e an intertwiner which, at each edge e , maps the $SO(4)$ -irreducible representations associated with the incoming 2-faces at the edge e to the outgoing $SO(4)$ -irreducible representations. $A_f(J_f)$ is a coefficient associated with each face f and so $A_e(j_f, i_e)$ and $A_v(j_f, i_e)$ with each edge e and vertex v respectively. One of the reason of it, we have found, is that the constraints (9) and (12), the delta function of the partition function (22), make the integrals highly non-local objects with some similarities with the measure of Quantum Regge Calculus [18]. It happens that the b^i 's, which are not adjacent and “very far”, can be correlated by implementing the constraints. This seems to prevent writing the partition functions (22) as a product of three factor as in (24). In BC-model, as well as other models for Spin Foam Formalism [14], the four-dimensional simplices, or equivalently, in the dual map, the dual vertices, are, in general, separated one from the others and there are no constraints to be implemented so the partition function results to be factorized as in (24).

IV. COUPLING WITH FERMIONIC MATTER

The coupling with fermionic matter, in the continuum case on Riemannian manifolds with torsion-free connection, is given by the lagrangian density

$$\mathcal{L} \equiv \frac{i}{2} \left(\bar{\psi} e_a^\mu \gamma^a \nabla_\mu \psi - e_a^\mu (\overline{\nabla_\mu \psi}) \gamma^a \psi \right) - m \bar{\psi} \psi \quad . \quad (25)$$

The γ^a , $a = 1, \dots, n$ are the Dirac-matrices satisfying the Clifford algebra

$$\gamma^a \gamma^b + \gamma^b \gamma^a = 2\delta^{ab} \quad , \quad (26)$$

whereas ψ is the n -dimensional Dirac spinor field ($\bar{\psi} \equiv \psi^\dagger \gamma^1$), ∇_μ the covariant derivative, and e_a^μ the n -beins on the tangent space of the Riemannian manifold (M, g) .

Now we have all the ingredients to define the coupling of gravity with fermionic matter on the lattice in analogy with the continuum case[19]. Let $2\nu = n$ or $n = 2\nu + 1$ (depending on whether n is even or odd) and consider the 2^ν -dimensional representation of the two-fold covering group of $SO(n)$ [10]. So, instead of considering the connection matrices $\Lambda_{\alpha\beta}$, we will deal with the $2^\nu \times 2^\nu$ connection matrices $D_{\alpha\beta}$ such that

$$D_{\alpha\beta} \gamma^a D_{\alpha\beta}^{-1} = (\Lambda_{\alpha\beta})_b^a \gamma^b \quad . \quad (27)$$

Given $D_{\alpha\beta}$ we can determine $\Lambda_{\alpha\beta}$. Furthermore if we know $\Lambda_{\alpha\beta}$, we can determine $D_{\alpha\beta}$ up to a sign. In particular, from (27), we can write $\Lambda_{\alpha\beta}$ as [10]

$$(\Lambda_{\alpha\beta})_{ab} = \frac{1}{2^\nu} \text{Tr}(\gamma_a D_{\alpha\beta} \gamma_b D_{\alpha\beta}^{-1}) \quad (28)$$

In the discrete theory we assume that the spinor field is a 2^ν complex vector defined at each vertex of the dual Voronoi complex, that is to say a map that to each vertex α associates the 2^ν complex vector $\psi(\alpha)$.

Let $|\alpha\beta|$ the distance between the two neighboring circumcenters in α and β [9]. We are ready to define the covariant derivative $(\nabla_\mu \psi)(\alpha)$ on a lattice

$$(\nabla_\mu \psi)(\alpha) \equiv \frac{D_{\alpha\beta} \psi(\beta) - \psi(\alpha)}{|\alpha\beta|} \quad . \quad (29)$$

So far the discrete version of the the action for the coupling between gravity and fermionic matter can be written in the form (see also [19])

$$\begin{aligned} S^F \equiv \sum_{\alpha} \left(\sum_{(\alpha\beta), \beta=1, \dots, n+1} \frac{i}{2|\alpha\beta|} \left(\overline{\psi(\alpha)} b_{\alpha\beta}^a \gamma_a D_{\alpha\beta} \psi(\beta) \right. \right. \\ \left. \left. - \left(\overline{D_{\alpha\beta} \psi(\beta)} \right) b_{\alpha\beta}^a \gamma_a \psi(\alpha) \right) - mV(\alpha) \overline{\psi(\alpha)} \psi(\alpha) \right) \quad . \quad (30) \end{aligned}$$

Then the quantum measure, which also includes fermionic matter, can be written as

$$\begin{aligned}
Z = \int e^{-(S+S^F)} \prod_{\alpha} \mu(\psi(\alpha)) \mu(\overline{\psi(\alpha)}) \delta\left(\sum_{\beta=1}^{n+1} b_{\alpha\beta}(\alpha)\right) \\
\times \prod_{\alpha\beta} \delta(b_{\alpha\beta}(\alpha) - D_{\alpha\beta} b_{\beta\alpha}(\beta)) \mu(D_{\alpha\beta}) \mu(b_{\alpha\beta})
\end{aligned} \tag{31}$$

where S is the action for pure gravity (14), $\mu(D_{\alpha\beta})$ the Haar measure on the two-fold covering group of $SO(n)$, while $\mu(\psi(\alpha)) = d\psi(\alpha)$ and $\mu(\overline{\psi(\alpha)}) = d\overline{\psi(\alpha)}$ are Grassmann variables.

Now on let's focus our reasonings in four dimensions. The two-fold covering group of $SO(4)$ is $Spin(4) = SU(2) \times SU(2)$. If we denote with

$$S_{\alpha\beta}^F \equiv \frac{i}{2|\alpha\beta|} \left(\overline{\psi(\alpha)} b_{\alpha\beta}^a \gamma_a D_{\alpha\beta} \psi(\beta) - \left(\overline{D_{\alpha\beta} \psi(\beta)} \right) b_{\alpha\beta}^a \gamma_a \psi(\alpha) \right) ,$$

and with

$$S_{\alpha}^F \equiv -mV(\alpha) \overline{\psi(\alpha)} \psi(\alpha) , \tag{32}$$

we can expand $e^{S_{\alpha\beta}^F}$ in Fourier series on the compact group $Spin(4)$ [13]

$$e^{-S_{\alpha\beta}^F} = \sum_I \sum_{ij} c_I^{ij}{}_{\alpha\beta}(\psi, \overline{\psi}, b_{\alpha\beta}) D_{\alpha\beta}^I{}_{ij} \tag{33}$$

where

$$c_I^{ij}{}_{\alpha\beta}(\psi, \overline{\psi}, b_{\alpha\beta}) \equiv c_I^{ij}{}_{\alpha\beta}(\psi(\alpha), \overline{\psi(\alpha)}, \psi(\beta), \overline{\psi(\beta)}, b_{\alpha\beta}) \equiv d_I \int_{Spin(4)} e^{-S_{\alpha\beta}^F} (\overline{D_{\alpha\beta}})^I{}_{ij} \mu(D_{\alpha\beta}) , \tag{34}$$

with I denoting the irreducible representations of $Spin(4)$ and d_I the dimension of the irreducible representation. Then we can rewrite equation (31) in the following way:

$$\begin{aligned}
Z = \prod_{\alpha\beta} \sum_I \sum_{ij} c_I^{ij}{}_{\alpha\beta}(\psi, \overline{\psi}, b_{\alpha\beta}) D_{\alpha\beta}^I{}_{ij} \delta(b_{\alpha\beta}(\alpha) - D_{\alpha\beta} b_{\beta\alpha}(\beta)) \mu(D_{\alpha\beta}) \mu(b_{\alpha\beta}) \\
\prod_{h \supset \alpha\beta} \sum_J c_J^h \left(b_{\alpha\beta}, b_{\beta\alpha}, \dots, b_{\delta_k^h \alpha}, b_{\alpha \delta_k^h} \right) \chi_J(D^h) \prod_{\alpha} e^{-S_{\alpha}} \delta\left(\sum_{\beta=1}^{n+1} b_{\alpha\beta}(\alpha)\right) \mu(\psi(\alpha)) \mu(\overline{\psi(\alpha)}) , \tag{35}
\end{aligned}$$

in which $\chi_J(D^h)$, according to equation (20), is the character of the J th-irreducible representation of the two-fold covering D^h of the holonomy matrix U^h around the plaquette h .

Although this partition function seems pretty complicated, it allows a straightforward *combinatorial* interpretation. If, as it seems very plausible even if not proved yet, Local Regge Calculus is related to Spin Foam Formalism then the presence of two independent representation indices I and J , in the path integral (35), can be interpreted, from a diagrammatical and combinatorial point of view as well, as we can associated each irreducible representation J with the **interior** of each plaquette h and each irreducible representation I with each dual-edge $\alpha\beta$. This corresponds, graphically, to color the interior of each plaquette and each edge with, in general, **different** colors! This would imply that the correspondings Spin Networks, as projection along the evolution direction [14] of Spin Foam 2-complex, of gravity with fermionic matter would be characterized by the fact that a $Spin(4)$ representation J is associated to each edge and, pairwise, an independent, from J , $Spin(4)$ representation I is associated to each vertex as well.

V. CONCLUSIONS

We have introduced a discrete theory of gravity in its first-order formalism. We have defined an action as the sum over the hinges of the traces of the holonomy matrices multiplied by the oriented volumes of the hinges. This action can be written on the dual Voronoi complex of the original simplicial complex. We can express the action as a function of the connection matrices and of the vectors $b_{\alpha\beta}$ which are normal to the faces of the simplices and whose modulo is proportional to the volume of the faces themselves.

The action is very similar to the Wilson action of lattice gauge theory[20]. Here the $b_{\alpha\beta}$ have the same role as the n -bein in the continuum theory. Moreover the action is locally invariant under the action of Lorentz group.

A quantum measure and the relative partition function has been defined. They are locally invariant under the action of $SO(n)$. The expansion of the exponential factor, in the partition function, shows that there are some similarities between Local Regge Calculus and Spin Foam Formalism. However many severe difference seems to show up as well, mainly based on the fact that, due to the presence of constraints on the dynamical variable which have to be implemented at quantum level, the quantum measure of Local Regge Calculus, as in the case of Regge Calculus [18], is *highly* non local. This generates differences between the partition function of Local Regge Calculus and Spin Foam Formalism. We guess that, since

the action of Local Regge Calculus converges in *the continuum limit* to the Einstein-Hilbert theory, whereas none have proved yet that Spin Foam Formalism converges to Gravity and Spin Foam Formalism is mainly based on many different models, a possible future work would consist of showing how we can derive different models used in Spin Foam Formalism from Local Regge Calculus doing suitable approximations. This should demonstrate that Local Regge Calculus contains Spin Foam Formalism. We have introduced a coupling of fermionic matter with discrete gravity. This coupling seems very natural and might be, in case, a way to introduce Fermionic matter and Gauge Fields as well in Spin Foam Formalism. As in the *pure-Gravity* case, the expansion of the exponential factor shows strict analogies with Spin Foam Formalism but, we have seen, the introduction of fermionic matter implies some peculiar labeling of the 2-complex which could affect, pairwise, the Spin Foam Complex and the relative Spin Network as well.

Acknowledgments

I would like to thank Alessandro D'Adda for the early and determinant collaboration on first-order Regge Calculus, Hugo A. Morales Tecotl for discussion and Carlo Rovelli for collaboration on the last part of this paper. I would like to express my gratitude to Fr. Bill Stoeger S.J. for directing, encouraging, and advising me. It is a pleasure to thank Fr. George Coyne S.J. and Fr. Francesco Tata S.J. for contributions to this research.

-
- [1] T. Regge, *General relativity without coordinates*, Nuovo Cimento **4**, Vol. 19 pag. 559-571 (1961)
 - [2] T. Regge, R.M. Williams, *Discrete structures in gravity*, Jour. of Math. Phys. **6**, Vol. 41 pag. 3964-3984 (2000).
 - [3] H. Seifert and W. Threlfall, *Seifert and Threlfall: a Textbook of Topology* (Academic Press, 1980).
 - [4] W.P. Thurston, *Shapes of polyhedra and triangulations of the sphere*, math.GT/9801088 (1998).
 - [5] H. W. Hamber, *Simplicial quantum gravity*, in K. Osterwalder and R. Stora eds., Critical phenomena, random systems, gauge theories, Les Houches Session XLII 1984 (Elsevier Science Publishers B.V., 1986).

- [6] M. Caselle, A. D’Adda, L. Magnea, *Regge Calculus as a local theory of the Poincaré group*, Phys. Lett. B **4**, Vol. 232 pag. 457-461 (1989).
- [7] M. Senechal, *Quasicrystal and geometry* (Cambridge University Press 1995).
- [8] G. Gionti, *Discrete Approaches Toward the Definition of a Quantum Theory of Gravity*, gr-qc/9812080, Ph.D. thesis SISSA (1998).
- [9] G. Gionti, *Discrete Gravity as a local theory of the Poincaré group*, Class. Quantum Grav. **22**, pag. 4217-4232 (2005).
- [10] J. Fröhlich, *Regge calculus and discretized gravitational functional integrals*, Non-perturbative quantum field theory -mathematical aspects and applications (World Scientific, Singapore 1992).
- [11] J. Cheeger, W. Muller, R. Schrader, *On the curvature of Piecewise flat manifolds*, Commun. Math. Phys., Vol. 92 pag. 405-454 (1984).
- [12] G. Esposito, G. Gionti, C. Stornaiolo, *Space-Time Covariant Form of Ashtekar’s Constraints*, Nuovo Cimento B **110**, pag. 1137-1152 (1995).
- [13] N.J. Vilenkin, *Special Function and the Theory of Group Representations* (Providence, Rhode Island: American Mathematical Society, 1968).
- [14] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004)
- [15] D. Oriti, *Spin Foam Models of Quantum Space-Time*, gr-qc/0311066, Ph.D. thesis DAMTP (2003).
- [16] J.W. Barrett, L. Crane, *Relativistic Spin Networks and Quantum Gravity*, J. Math. Phys. **29**, pag. 3296-3302 (1998).
- [17] M.P. Reisenberger, *Classical Euclidean general relativity from “left-handed area=right-handed area”*, gr-qc/9804061 (1998).
- [18] P. Menotti, P.P. Peirano, *Functional Integration on Two-Dimensional Regge Geometries*, Nucl. Phys. B, Vol. 473, pag. 426-454 (1996).
- [19] H. Ren, *Matter Fields in Lattice Gravity*, Nucl. Phys. B, Vol. 301 pag. 661-684 (1988).
- [20] K. T. Wilson, *Confinement of Quarks*, Physical Review D **10**, pag. 2445-2459 (1974).